

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/040513

International filing date: 03 December 2004 (03.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/528,074  
Filing date: 05 December 2003 (05.12.2003)

Date of receipt at the International Bureau: 31 January 2005 (31.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1275101

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

*January 19, 2005*

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.**

**APPLICATION NUMBER: 60/528,074**

**FILING DATE: *December 05, 2003***

**RELATED PCT APPLICATION NUMBER: *PCT/US04/40513***



Certified by

Under Secretary of Commerce  
for Intellectual Property  
and Director of the United States  
Patent and Trademark Office

Express Mail Label No.: EV312715232US

Date of Deposit: December 5, 2003

Attorney Docket No. 19374-515 PRO

**PROVISIONAL APPLICATION FOR PATENT COVER SHEET**

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR § 1.53(c)

15535 U.S. PTO  
601528074



INVENTORS / APPLICANTS		
Given Name (last name, first name, and middle initial [if any])		Residence (City and State or Foreign Country)
Farmer Lefkowitz	Sean Andrew	R. 1520 Stillwater Drive, Miami Beach, FL 33141 24400 Highpoint Road, Suite 5, Beachwood, Ohio 44122
TITLE OF THE INVENTION		
METHODS FOR THE DIETARY MANAGEMENT OF IRRITABLE BOWEL SYNDROME AND CARBOHYDRATE MALABSORPTION		
CORRESPONDENCE ADDRESS		
Attorney Name: Firm Name and Address:	Ivor R. Elrifi, Ph.D. MINTZ, LEVIN, COHN, FERRIS, GLOVSKY AND POPEO, P.C. One Financial Center Boston, MA 02111	
Telephone:	(617) 542-6000	
Fax:	(617) 542-2241	
ENCLOSED APPLICATION PARTS		
<input checked="" type="checkbox"/> Specification	Number of Pages: 31	
<input type="checkbox"/> Sequence Listing	Number of Pages:	
<input type="checkbox"/> Figures	Number of Sheets:	
<input type="checkbox"/> Other documents (specify):		
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government:		
<input checked="" type="checkbox"/> No.		
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are:		
METHOD OF PAYMENT		
<input checked="" type="checkbox"/> A check in the amount of \$80.00 is enclosed to cover the filing fees of the Provisional application.		
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge \$80.00 to cover the filing fees of the Provisional application to Deposit Account No. 50-0311, Reference No. 19374-515.		
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge additional fees or credit any overpayment to Deposit Account No. 50-0311, Reference No. 19374-515.		

Respectfully submitted,

*for*   
Ivor R. Elrifi, Reg. No. 39,529  
Ingrid A. Beattie, Reg. No. 42,306  
c/o MINTZ, LEVIN  
Tel: (617) 542-6000  
Fax: (617) 542-2241  
Customer No.: 30623

Dated: December 5, 2003

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

SEND TO: MAIL STOP PROVISIONAL APPLICATION, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450

**METHODS FOR THE DIETARY MANAGEMENT OF IRRITABLE BOWEL  
SYNDROME AND CARBOHYDRATE MALABSORPTION**

**TECHNICAL FIELD OF THE INVENTION**

The present invention relates to the utilization of probiotic lactic acid-producing bacteria in a nutritional composition. More specifically, the present invention relates to the use of *Bacillus coagulans* for increasing the absorption of carbohydrates within the gastrointestinal tract of a mammal.

**BACKGROUND OF THE INVENTION**

The human digestive system uses a series of enzymes to break down complex foods into simple molecules (*e.g.*, sugars, peptides and lipids) that can be absorbed by the body. The inability or diminished capacity of the body's production of one or more enzymes that are crucial for proper digestion by can lead to gastrointestinal symptoms that have been characterized by the medical community as irritable bowel syndrome (IBS). A patient with IBS will present clinically with one of three variants: i) chronic abdominal pain and constipation (also known as spastic colitis); ii) chronic intermittent diarrhea, often without pain; or iii) both features, in an alternating cycle of constipation and diarrhea.

**SUMMARY OF THE INVENTION**

The invention is based, in part, on the discovery of the therapeutic effects of *Bacillus coagulans*, a spore-forming lactic acid bacterium, in the prevention and treatment of IBS and carbohydrate malabsorption. Carbohydrate malabsorption includes the inability of a mammal to fully digest the naturally occurring sugars (*e.g.*, lactose, fructose, and glucose) in foods and beverages.

In one aspect, the invention provides a method of reducing one or more symptoms of irritable bowel syndrome, by identifying a patient suffering from or at risk of developing irritable bowel syndrome, and administering to the patient a composition that includes *Bacillus coagulans* bacteria. Bacterial species include *Bacillus coagulans*, *e.g.*, *Bacillus*

*coagulans* hammer, preferably *Bacillus coagulans* hammer strain Accession No. ATCC 31284, or strains derived from *Bacillus coagulans* hammer strain Accession No. ATCC 31284. Symptoms of IBS include diarrhea, constipation, alternating diarrhea and constipation, bloating, urgency, and abdominal pain. In embodiments of the invention, the composition also includes a supplementary enzyme (*e.g.*, a lactase, a fructase, a lipase, or a peptidase), an anti-diarrheal agent (*e.g.*, loperamide, attapulgit, Croton Lechleri Extract, or calcium polycarbophil), or a laxative (*e.g.*, a senoside such as senosides A, B, C or D, docusate sodium, magnesium hydroxide, or a dietary fiber). A supplemental lactase includes an enzyme that is capable of catalyzing the hydrolysis of lactose in the gastrointestinal tract of a mammal, in a concentration that exceeds the amount of lactase that is available in the small and large intestines of a mammal prior to *Bacillus coagulans* colonization. A supplemental fructase includes an enzyme that is capable of catalyzing the hydrolysis of fructose in the gastrointestinal tract of a mammal, in a concentration that exceeds the amount of fructose that is available in the small and large intestines of a mammal prior to *Bacillus coagulans* colonization. The gastrointestinal tract is the system of organs in a mammal including the mouth (buccal cavity), pharynx, esophagus and cardia, stomach(s), and intestines.

Colonization of *Bacillus coagulans* bacteria generally occurs between 24-48 hours following delivery. Continued colonization is improved by the repeated administration of *Bacillus coagulans*, such as daily administration. Generally, the supplementary fructase is provided at a dose of from about 1000IU to about 12,000IU, and the supplementary lactase is provided at a dose of from about 1000IU to about 12,000IU. In embodiments of the invention, the patient is female, such as a female that is post-menstrual or post-menopausal. Alternatively, the patient is male.

The invention provides *Bacillus coagulans* bacteria at a concentration of from about  $1 \times 10^4$  to about  $1 \times 10^{12}$  viable bacteria, specifically about  $1 \times 10^6$  to about  $1 \times 10^{11}$ , more specifically about  $1 \times 10^8$  to about  $1 \times 10^{10}$ , and most specifically about  $8 \times 10^8$ . The *Bacillus coagulans* bacteria are in the form of spores, vegetative cells, or a combination thereof.

The invention also provides a method of reducing one or more symptoms of irritable bowel syndrome, by identifying a patient suffering from or at risk of developing irritable

bowel syndrome, and administering to said patient a composition including an effective IBS inhibiting amount of *Bacillus coagulans* bacteria prior to or concomitant with the onset of one or more IBS symptoms. Symptoms of IBS include diarrhea, constipation, alternating diarrhea and constipation, bloating, urgency, and abdominal pain.

5           In another aspect, the invention, the invention provides a method of reducing a symptom of irritable bowel syndrome, by identifying a patient suffering from or at risk of developing irritable bowel syndrome, and administering to the patient a composition that includes an enzyme, preferably a fructase and a lactase.

10           The invention also provides a method of diagnosing irritable bowel syndrome in a patient, including the steps of identifying a patient having a symptom of irritable bowel syndrome, providing a patient-derived biological sample from the identified patient, determining an amount of a product of a gastrointestinal enzyme in the patient-derived sample, and comparing the amount in the patient-derived sample with a reference amount of a product of a gastrointestinal enzyme, whereby an alteration in the test amount relative to  
15           the reference amount indicates that the patient has irritable bowel syndrome.

          A gastrointestinal enzyme includes any enzyme that is active in the gastrointestinal tract, particularly the stomach and the small and large intestines. A biological sample includes any solid, liquid, or gaseous material obtained from a mammal, such as a human patient. The symptoms of IBS include diarrhea, constipation, or alternating diarrhea and  
20           constipation. The gastrointestinal enzyme includes a lactase, a fructase, a lipase and a peptidase. In embodiments of the invention, the patient identified as having a symptom of irritable bowel syndrome has one or more symptoms classified under the Rome Criteria. The amount of the product of a gastrointestinal enzyme in the patient-derived sample is modulated following administration of the compositions of the invention. For example,  
25           hydrogen measured using the hydrogen breath test declines following administration of a *Bacillus coagulans*-containing composition.

          In another aspect, the invention provides a method of improving stool consistency in a patient afflicted with non-constipated IBS, by administering an effective amount of a *Bacillus coagulans* bacteria provided at a concentration of from about  $1 \times 10^8$  to about  $1 \times 10^{10}$

viable bacteria, where the patient's stool consistency is improved following the administration.

In a further aspect, the invention relates to a method of decreasing urgency in a subject afflicted with IBS, by administering an effective amount of a *Bacillus coagulans* bacteria provided at a concentration of from about  $1 \times 10^8$  to about  $1 \times 10^{10}$  viable bacteria wherein urgency is decreased following the administration.

The invention also provides a composition that includes a *Bacillus coagulans* bacteria, a supplementary lactase (e.g.,  $\beta$ -galactosidase), and a supplementary fructase. Generally, the supplementary lactase is provided in a concentration from about 1000 IU to about 12,000 IU (e.g., about 3000IU), and the supplementary fructase is provided in a concentration from about 1000 IU to about 12,000 IU (e.g., about 3000IU). In embodiments of the invention, the composition also includes an anti-diarrheal agent, a laxative, a vitamin, a mineral, an isolated amino acid, a source of dietary fiber, or an antibiotic. The composition may also include a pharmaceutically-acceptable carrier containing, e.g., silicone. The composition is provided in the form of a capsule, tablet (including a chewable tablet), powder, liquid or in a formulation with a food product. Food products include dairy products including ice cream, nutritional bars (energy or candy bars), sugar substitutes, non-dairy creamers, tea bags, and similar products. Sources of dietary fiber include psyllium husk, soy fiber, citrus fiber, beet fiber, pumpkin seed meal, ground flax, black walnut hull, rice fiber, hydrocollodial polysaccharides, pecan husks, and peanut husks.

The invention provides a composition comprising a *Bacillus coagulans* bacteria and a supplemental enzyme provided in a formulation with a food product. In embodiments of the invention, the food product is a dairy product (a product containing a component obtained from the milk of a cow, sheep, goat, or similar mammal).

The invention further provides a composition that includes from about  $1 \times 10^8$  to about  $1 \times 10^{10}$  *Bacillus coagulans* bacteria, a supplemental lactase provided in a concentration of about 3000 IU, a supplemental fructase provided in a concentration of about 3000 IU, and manganese stearate.

The invention also provides a composition that includes an isolated lactase provided in a concentration from about 1000 IU to about 12,000 IU per dose and an isolated fructase provided in a concentration from about 1000 IU to about 12,000 IU per dose. In embodiments of the invention, the composition also includes an anti-diarrheal agent, a  
5 laxative, a vitamin, a mineral, an isolated amino acid, a source of dietary fiber, an antibiotic, or a combination thereof.

The invention also provides a method for increasing carbohydrate absorption in a mammal, by administering to a mammal a composition that includes a *Bacillus coagulans* bacteria, a supplementary lactase (*e.g.*, provided in a concentration from about 1000 IU to  
10 about 12,000 IU), and a supplementary fructase (*e.g.*, provided in a concentration from about 1000 IU to about 12,000 IU), where carbohydrate absorption is increased following the administration. In embodiments, the mammal is diagnosed as suffering from or being at risk of developing a disorder associated with carbohydrate malabsorption. Disorders associated with carbohydrate malabsorption include lactose intolerance, fructose intolerance, glucose-  
15 galactose intolerance, sorbitol intolerance, irritable bowel syndrome, short bowel syndrome, stagnant loop syndrome, celiac disease, chronic malnutrition, chronic persistent diarrhea, immunoproliferative small intestinal disease, intractable diarrhea of infancy, postenteritis syndrome, tropical sprue, Whipple's disease, Wolman disease, Crohn's disease and ulcerative colitis. In embodiments, the composition also includes an anti-diarrheal agent, a laxative, a  
20 vitamin, a mineral, an isolated amino acid, a source of dietary fiber, or an antibiotic.

The invention further provides a method for increasing lactose digestion, including the steps of identifying a patient suffering from or at risk of developing lactose intolerance, and administering to the patient a composition that includes *Bacillus coagulans* bacteria and a supplemental lactase (*e.g.*, provided in a concentration from about 1000 IU to about 12,000  
25 IU), whereby lactose digestion is increased following the administration.

The invention also provides a composition including a *Bacillus coagulans* bacteria and a supplementary fructase, *e.g.*, a fructase provided in a concentration from about 1000 IU to about 12,000 IU. In embodiments of the invention, the composition also includes an isolated amino acid. The composition is provided in the form of a capsule, tablet (including  
30 chewable tablet), powder, liquid or in a formulation with a food product. The *Bacillus*



*coagulans* bacteria are derived from *Bacillus coagulans* hammer strain Accession No. ATCC 31284.

In another aspect, the invention provides a medical food for the management of irritable bowel syndrome, that includes *Bacillus coagulans* bacteria and an isolated amino acid (e.g., L-lysine), wherein said medical food is formulated to provide at least about  $1 \times 10^6$  (e.g.,  $1 \times 10^7$ ,  $1 \times 10^8$  or  $8 \times 10^8$  or more) viable *Bacillus coagulans* bacteria in the gastrointestinal tract of a mammal per day based on a serving size of about 0.5 gram to about 25 grams of the medical food taken up to twice a day. In embodiments of the invention, the medical food is provided at a dosage such that colonization of about  $1 \times 10^5$  (e.g.,  $1 \times 10^6$  or  $1 \times 10^7$ ) viable *Bacillus coagulans* bacteria per gram of fecal material in a mammal following consumption of the medical food. In embodiments of the invention, the medical food includes a supplemental enzyme such as a lactase, a fructase, a lipase or a peptidase. In other embodiments, the medical food includes an anti-diarrheal agent, a laxative, a vitamin, a mineral, an appropriate amino acid(s), a source of dietary fiber, and/or an antibiotic.

In a further aspect, the invention provides a method of dietary management of a subject's carbohydrate absorption, including the steps of identifying a patient having a symptom of carbohydrate malabsorption, and providing a composition comprising *Bacillus coagulans* bacteria to the identified patient, wherein the bacteria colonize the subject's gastrointestinal tract, whereby carbohydrate absorption by the subject is modulated, such that the subject's carbohydrate absorption and is managed. The dietary management of the subject's carbohydrate absorption results in a reduction or elimination of one or more of the symptoms of carbohydrate malabsorption.

In another aspect, the invention provides a method of dietary management of a subject's carbohydrate absorption, including the steps of identifying a patient having a symptom of carbohydrate malabsorption, providing a patient-derived biological sample from the identified patient, determining an amount of a product of a gastrointestinal enzyme in the patient-derived sample, comparing the amount in the patient-derived sample with a reference amount of a product of a gastrointestinal enzyme, and providing a composition comprising *Bacillus coagulans* bacteria, whereby the subject's carbohydrate absorption is managed. In embodiments of the invention, the amount of a product of a gastrointestinal enzyme in the patient-derived sample is determined using hydrogen and/or methane breath testing. The

amount of the product in the patient-derived sample declines following administration of the *Bacillus coagulans*-containing composition. In further embodiments of the invention, the patient's carbohydrate absorption is managed such that one or more symptoms of carbohydrate malabsorption are decreased or eliminated.

5           The invention further provides a method for increasing carbohydrate absorption in a patient diagnosed as suffering from or being at risk of developing celiac disease, by administering to the subject a composition comprising *Bacillus coagulans* bacteria, wherein carbohydrate absorption in the patient is increased following the administration.

10           In another aspect, the invention provides a method of reducing a symptom of irritable bowel syndrome, wherein the symptom comprises alternating diarrhea and constipation, comprising identifying a patient suffering from or at risk of developing irritable bowel syndrome, and administering to the patient a composition including *Bacillus coagulans* bacteria.

15           Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, 20 and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

## DETAILED DESCRIPTION

25           Medical diagnosis of IBS was based on the absence or presence of a number of symptoms, which are generally regarded as typical of IBS and are provided, for example, by the "Rome Criteria" (See, W. G. Thompson et al., Gastroent. Int. 2 (1989) 92-95; W. G. Thompson et al., Gut 45/II (1999) II43-II 47; W. G. Thompson, Lancet 341 (1993) 1569-1572), and the Rome II Criteria. Guidelines for IBS diagnosis under the Rome criteria 30 include the continuous or recurrent symptoms of abdominal pain or discomfort that may be

relieved with defecation, may be associated with a change in frequency, or may be associated with a change in the consistency of stools; and that two or more of the following symptoms are present at least 25 percent of the time: altered stool frequency (greater than 3 bowel movement per day or less than 3 bowel movements per week), altered stool form (hard or loose watery stools or poorly formed stools), passage of mucous stools, and bloating (feeling of abdominal distention). The importance of carbohydrates in the onset of IBS symptoms has recently been discussed. (See, *Scand J Gastroenterol.* 1998. 33(11):1158-63, *Isr Med Assoc J.* 2000. 2(8):583-7, *Am J Gastroenterol.* 2003. 98(6):1348-53). IBS and carbohydrate malabsorption have often been confused for one another. Reliable diagnosis is critical to determining the appropriate dosage for treating the symptoms of IBS. The present invention provides diagnostic methods for the detection and diagnosis of IBS in patients, such as humans, who are suffering from IBS, or are at risk of developing IBS.

The present invention also provides methods for the reduction of symptoms of IBS. Prior to the present invention, it has been difficult to effectively reduce symptoms in the treatment of irritable bowel syndrome. The goals of therapeutic treatment were to reduce the variety of complaints, and to improve conditions so as to decrease morbidity and increase the quality of the patient's daily life. Therapeutic treatments include psychotherapy, life guidance, diet therapy, and drug therapy used on a symptomatic basis against the patient's complaints. Compounds including opioid agonists (e.g., loperamide) or anticholinergic agents (e.g., mepenzolate bromide and timentidine bromide) have been used to control hypermotility of the digestive tract, and benzodiazepine drugs (e.g., diazepam) have been prescribed for anxiety, insomnia and similar complaints. Recently, antagonists of 5-hydroxytryptamine (5-HT; serotonin) and 5-HT receptors have been used to treat IBS. (See, U.S. Patent 6,429,209). A deficiency of these treatments is that they are usually incapable of reducing or eliminating multiple symptoms of IBS, particularly when the patient presents with alternations of diarrhea and constipation. The compositions of the present invention alleviate multiple symptoms of carbohydrate malabsorption, including pain, flatus, abdominal bloating, diarrhea, constipation, and alternating diarrhea and constipation

The present invention also provides methods of treatment of diseases associated with carbohydrate malabsorption. These diseases, in addition to IBS, include lactose intolerance, fructose intolerance, glucose-galactose intolerance, sorbitol intolerance, short bowel

syndrome, stagnant loop syndrome, celiac disease, chronic malnutrition, chronic persistent diarrhea, immunoproliferative small intestinal disease, intractable diarrhea of infancy, postenteritis syndrome, tropical sprue, Whipple's disease, Wolman disease, Crohn's disease and ulcerative colitis.

## 5 *Bacillus coagulans*

*Bacillus coagulans* is a strain of bacteria that possesses the ability to sporulate, making the strain resistant to heat and other conditions, as well as providing for a long shelf-life in product formulations. Further, *Bacillus coagulans* is ideal for survival and colonization of tissues under conditions of pH, salinity, and the like within the  
10 gastrointestinal tract. Additionally, *Bacillus coagulans* is non-pathogenic. Preferred methods disclosed herein utilize *Bacillus coagulans* cells and spores. Methods of preparing *Bacillus coagulans* vegetative cells and spores are presented in Example 1.

The *Bacillus coagulans* bacteria of the present invention are capable of colonizing the gastrointestinal tract of a mammal to which they are provided enterically. Generally,  
15 colonization occurs within twenty four to forty eight hours following administration. Efficiency of intestinal colonization of a mammal can be determined, *e.g.*, by quantitating the number of *Bacillus coagulans* bacteria per gram of the mammal's feces. As used herein, a mammal has been colonized by the *Bacillus coagulans* bacteria of the present invention if the mammal's feces contain greater than  $1 \times 10^4$  viable bacteria per gram of feces, preferably  
20  $1 \times 10^5$  viable bacteria per gram of feces, or more preferably  $1 \times 10^6$  viable bacteria per gram of feces.

One species of *Bacillus coagulans*, that is useful in this invention, had previously been mischaracterized as a *Lactobacillus* in view of the fact that, as originally described, this bacterium was labeled as *Lactobacillus sporogenes*. However, initial classification was  
25 incorrect due to the fact that *Bacillus coagulans* produces spores and through metabolism excretes L(+)-lactic acid, both aspects which provide key features to its utility. Instead, these developmental and metabolic aspects required that the bacterium be classified as a lactic acid *Bacillus*, and therefore it was re-designated. Accordingly, the bacteria useful in the invention (i) possess the ability to produce and excrete enzymes useful in digestion (*e.g.*, lactase,  
30 various proteases, lipases and amylases); (ii) demonstrate beneficial function within the gastrointestinal tract; and (iii) are non-pathogenic.

The Gram positive rods of *Bacillus coagulans* have a cell diameter of greater than 1.0  $\mu\text{m}$  with variable swelling of the sporangium, without parasporal crystal production.

*Bacillus coagulans* is a non-pathogenic, Gram positive, spore-forming bacteria that produces L(+) lactic acid (dextrorotatory) under homo-fermentation conditions. It has been isolated from natural sources, such as heat-treated soil samples inoculated into nutrient medium (see *e.g.*, *Bergey's Manual of Systemic Bacteriology*, Vol. 2, Sneath, P.H.A. *et al.*, eds., Williams & Wilkins, Baltimore, MD, 1986). Purified *Bacillus coagulans* strains have served as a source of enzymes including endonucleases (*e.g.*, U.S. Pat. No. 5,200,336); amylase (U.S. Pat. No. 4,980,180); lactase (U.S. Pat. No. 4,323,651) and cyclo-malto-dextrin glucanotransferase (U.S. Pat. No. 5,102,800). In particular, *Bacillus coagulans* strains have been used as general nutritional supplements and agents to control constipation and diarrhea in humans and animals.

Various *Bacillus coagulans* bacterial strains which are currently commercially available from the American Type Culture Collection (ATCC, Manassas, VA) include the following accession numbers: *Bacillus coagulans* Hammer NRS 727 (ATCC No. 11014); *Bacillus coagulans* Hammer strain C (ATCC No. 11369); *Bacillus coagulans* Hammer (ATCC No. 31284); and *Bacillus coagulans* Hammer NCA 4259 (ATCC No. 15949). Purified *Bacillus coagulans* bacteria are also available from the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (Braunschweig, Germany) using the following accession numbers: *Bacillus coagulans* Hammer 1915 (DSM No. 2356); *Bacillus coagulans* Hammer 1915 (DSM No. 2383, corresponds to ATCC No. 11014); *Bacillus coagulans* Hammer (DSM No. 2384, corresponds to ATCC No. 11369); and *Bacillus coagulans* Hammer (DSM No. 2385, corresponds to ATCC No. 15949). *Bacillus coagulans* bacteria can also be obtained from commercial suppliers such as K.K. Fermentation (Kyoto, Japan) and Nebraska Cultures (Walnut Creek, CA). The present invention also includes strains or variants derived from *Bacillus coagulans* Hammer strain ATCC No. 31284.

*Bacillus coagulans* bacteria are provided in amounts sufficient to colonize the gastrointestinal tract of a mammal. The invention provides *Bacillus coagulans* bacteria at a concentration of from about  $1 \times 10^4$  to about  $1 \times 10^{12}$  viable bacteria, specifically about  $1 \times 10^6$  to about  $1 \times 10^{11}$ , more specifically about  $1 \times 10^8$  to about  $1 \times 10^{10}$ , and most specifically about  $810^8$ . *Bacillus coagulans* bacteria are provided as vegetative cells, spores, or a combination thereof.

## ***Fructose***

The fructose of the invention is preferably an enzyme that is capable of catalyzing the hydrolysis of fructose in the gastrointestinal tract of a mammal. Fructose can be purified from a fungus such as *Aspergillus oryzae*. Fructose is also commercially available from  
5 Specialty Enzymes and Biochemicals (Chino, CA), Spectrum Chemicals (Los Angeles, CA), and Solvay Enzymes (Edison, NJ).

Fructose activity is measured *in vivo* using the hydrogen ion breath test. A patient who has abstained from carbohydrates for at least twelve hours is given a 33% fructose solution (50g per 150 ml of water), and end-expiratory breath samples are collected before  
10 (the baseline value) and every 15-30 minutes for four to six hours after sugar ingestion. Hydrogen (and other gases such as methane) breath concentrations are measured using gas chromatography. A person is defined as fructose intolerant if a rise of at least 3 parts per million (ppm) over three consecutive breath tests from the baseline value or a value over 20 ppm following sugar ingestion is observed.

## ***Lactase***

The lactase of the invention is preferably an enzyme that is capable of catalyzing the hydrolysis of lactose in the stomach and/or intestine. In certain embodiments, two lactases with different optimum pH ranges are used (*e.g.*, a first lactase that has an optimum pH range that encompasses pH 3.0 to about pH 6.0, and a second lactase that preferably has an  
20 optimum pH range that encompasses about pH 6.0 to about pH 8.0). An optimum pH range means the pH over which the hydrolytic activity of the lactase is within about 10 to 100 percent of its maximum, and optimum pH value means the pH at which the lactase exhibits maximum hydrolytic activity.

Lactases derived from fungi are generally known to have optimum pH values which  
25 fall within the acid range. Genera of fungi useful in obtaining lactases include *Aspergillus*; *Mucor*; *Fusarium*; *Scopulariopsis*; *Alternaria*; and *Curvularia* and the bacterium *Thermus aquaticus*. The lactases, having the optimum pH value shown in the parentheses, are preferably derived from the following fungi: *Aspergillus oryzae*; (4.5-5.0) *Aspergillus niger* (3.0-4.0); *Fusarium moniliforme* (3.8-5.0); *Scopulariopsis* (3.6-5.0); *Mucor pucillus* (4.5-6),  
30 *Alternaria alternata* (4.5-5.3); and *Curvularia inaequalis* (3.4-4.3) and the bacterium *Thermus aquaticus* (4.5-5.5).

Lactases derived from yeast and bacteria are generally known to have optimum pH values in the more neutral region, including *Kluyveromyces* (*Saccharomyces*), *Lactobacillus*, *Bacillus*, *Streptococcus*, and *Escherichia*. Lactase derived from the following organisms, having the optimum pH value shown in the parentheses, are preferred: *Kluyveromyces lactis* (6.5), *Kluyveromyces fragilis* (6.6), *Lactobacillus thermophilus* (6.2-7.1), *Bacillus circulans* (6.0), *Lactobacillus bulgaricus* (7.0), *Leuconostoc citrovorum* (56.5), *Bacillus stearo-thermophilus* (6.0-6.4), *Streptococcus thermophilus* (6.5-7.5), and *Bacillus* sp. (6.8).

The lactases used in the present invention can be produced by a variety of well known techniques. Many of these lactases are produced by commercial processes which cultivate the bacterium, yeast or fungus, and then isolate the lactase from the culture or culture broth of the microorganism. Further techniques for preparing such lactases may be found in U.S. Pat. No. 3, 629,073; U.S. Pat No. 3,718,739; and U.S. Pat. No. 3,919,049, all of which are hereby incorporated by reference.

Lactase activity is measured *in vivo* using the hydrogen ion breath test. A patient who has abstained from carbohydrates for at least twelve hours is given a lactose solution (18-50g), and end-expiratory breath samples are collected before and every 15-30 minutes for four to six hours after sugar ingestion. Hydrogen (and other gases such as methane) breath concentrations are measured using gas chromatography. An increase in breath hydrogen concentration of 10 parts per million (ppm) following sugar ingestion is typically observed in non-lactose intolerant patients. Incomplete absorption of carbohydrate is defined as an increase in breath hydrogen of 20 ppm (or its equivalent of 5 ppm in methane) following sugar ingestion.

Lactase activity is quantified in units. An FCC lactase unit (FCC Lac U), and IU and a neutral lactase unit are defined as that quantity of enzyme that will liberate 1  $\mu$ mol of o-nitrophenol from o- nitrophenyl- $\beta$ -D-galactoside per minute under the conditions, of the assay described in Food Chemicals Codex, National Academy Press, Wash., D. C., pp. 491-2 (1981), which is hereby incorporated by reference, at pH 4. 5 and 6.5, respectively.

#### ***Other gastrointestinal enzymes***

Amylase ( $\alpha$ -1,4-glucan 4-glucanohydrolase, EC 3.2.1.1.) activity is determined using the method of Somogyi (See, Somogyi, 1960. "Modification of two methods for the assay of amylase." *Clin Chem.* 6:23-35). One amylase activity unit is defined as the amount of

amylase that will cause the formation of reducing power equivalent to 1 mg glucose in 30 minutes at 40 degrees C per milligram of intestinal digesta protein. Cornstarch is an amylase substrate useful for calibration of amylase activity units.

Lipase (*e.g.*, LPS, triacylglycerol lipase, EC 3.1.1.3.) activity is assayed using the method of Tietz and Fiereck (See, Tietz and Fiereck, 1966. Clin. Chim. Acta 13:352-58). One lipase activity unit is equal to the volume (mL) of 0.05 M NaOH required to neutralize the fatty acid liberated during a 6 hour incubation with 3 mL of lipase substrate (*e.g.*, olive oil) at 37 degrees C per milligram of digesta protein. Lipases can be purified from *Bacillus subtilis* and *Pseudomonas aruginosa* and is commercially available.

Peptidases and proteases include enzymes that degrade a polypeptide by hydrolysis of the peptide bonds. Peptidases include amino-, dipeptidyl-, and tripeptidyl-peptidases. Useful proteases include Arg-C proteinase, Asp-N endopeptidase, caspases, chymotrypsin, clostripain, enterokinase, granzyme B, glutamyl endopeptidase, pepsin, proline-endopeptidase, proteinase K, Staphylococcal peptidase I, thermolysin, thrombin and trypsin.

### ***Composition Formulations***

The compositions of the present invention can be combined with a pharmaceutically acceptable carrier and administered orally. The unit dosages of these compositions may be in the form of solid preparations, such as tablets, pills, capsules, caplets, powders, granules and wafers, or liquid preparations, such as suspensions or dispersions in aqueous or non-aqueous vehicles, such as syrups and elixirs.

In some embodiments, the compositions of the present invention may contain components in addition to the *Bacillus coagulans* bacteria. Additional components include supplementary enzymes, anti-diarrheal agents, laxatives, dietary fibers, isolated amino acids, vitamins, minerals, antibiotics, and buffering agents.

Supplementary enzymes include a lactase, a fructase, a lipase, and a peptidase. Generally, a supplementary enzyme is provided in an amount exceeding the amount of the enzyme contained in the *Bacillus coagulans* bacteria provided. For example, a supplemental lactase is an amount of lactase sufficient to digest the lactose present in the gastrointestinal tract of a mammal when the supplementary lactase is consumed, in addition to any lactose



that is consumed prior to *Bacillus coagulans* colonization in the mammal's gastrointestinal tract.

Anti-diarrheal agents include any compounds that reduce diarrhea, such as by reducing water content in the stool. Preferred anti-diarrheal agents include loperamide (such as loperamide HCL), attapulgit, Croton Lechleri Extract, and calcium polycarbophil.

Laxatives include any compound that increases stool density or frequency of bowel movements. Preferred laxatives include senosides, docusate sodium, magnesium hydroxide, and a dietary fiber. Senosides include hydroxyanthracene glycosides such as senosides A, B, C or D, generally obtained from pulverized *Cassia acutifolia* husk. Exemplary dietary fibers include psyllium husk, soy fiber, citrus fiber, beet fiber, pumpkin seed meal, ground flax, black walnut hull, rice fiber, hydrocollodial polysaccharides, pecan husks, and peanut husks.

Isolated amino acids include alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine. Preferably, the isolated amino acid is lysine.

The compositions of the present invention are usable in the form of medicines, foods, and drinks, including supplements, medical foods, health foods, nutraceuticals, and dietary supplements, as directed by a healthcare practitioner. A medical food is prescribed by a physician when a patient has special nutrient needs in order to manage a disease or health condition such as IBS or carbohydrate malabsorption, and the patient is under the physician's ongoing care.

In preparing solid unit dosage forms, the compositions of the present invention (*e.g.*, *Bacillus coagulans* bacteria) are mixed with conventional solid fillers or carriers, such as silicone, starch, talc, calcium phosphate, calcium sulfate, calcium stearate, magnesium stearate, stearic acid, sorbitol, mannitol, gelatin, natural or synthetic gums, such as carboxymethylcellulose, methylcellulose, alginates, dextrans, acacia gum, karaya gum, locust bean gum, tragacanth and other conventional carriers. Additionally, other excipients such as diluents, binders, lubricants, disintegrants, colors and flavoring agents may be employed.

Suitable liquid forms of the present invention can be prepared by incorporating the lactase in aqueous or non-aqueous dispersions, suspensions, or solutions. Conventional liquid

carriers such as glycerol, and edible glycols, edible oils, such as cottonseed oil, soybean oil, corn oil, peanut oil, safflower oil, and other triglyceride oils, and dispersing or suspending agents, such as the aforementioned natural and synthetic gums.

Conventional methods are employed for preparing the solid and liquid forms of the present invention. Suitable techniques are described in *Remington's Pharmaceutical Sciences*, 18th Ed., Chapters 83 and 89 (1990), which is hereby incorporated by reference.

The compositions of the present invention can also be produced in powdered or granular form for direct admixture with food products consumed by subjects suffering from IBS or other carbohydrate malabsorption diseases. For instance, in the case of a lactose intolerant infant, a suitable amount of the *Bacillus coagulans* and lactase-containing composition of the present invention, in a powdered or granular form, can be added directly to the milk or other food consumed by the infant. In the case of an animal, such as a mammal, that normally requires a dietary regime of whey, the compositions of the present invention may be added directly to the whey.

In accordance with another embodiment of the present invention, the composition may contain an enteric coating, such as coating a *Bacillus coagulans* bacterium as a vegetative cell. This coating remains intact in the stomach, but will dissolve and release the vegetative cell once it reaches the more neutral environment of the small intestine. Suitable enteric coatings include amylose acetate phthalates, styrene-maleic acid copolymer, cellulose acetate succinate, cellulose acetate phthalate, polyvinyl acetate phthalate, hydroxy-propylmethylcellulose phthalate, fatty acids, fatty acid esters, glycerol esters, polyglycerol esters, paraffin waxes, carnauba wax, formalized gelatin, shellac and hydrogenated vegetable waxes, such as hydrogenated castor oil and cottonseed oil. Other suitable enteric coatings are disclosed in Lieberman, H. A. et al., *Pharmaceutical Dosage Forms: Tablets*, Vol. 3, pp. 114-116 (1990), which is hereby incorporated by reference. The enteric coating is applied using conventional particle coating techniques. If the vegetative cell is granulated with other excipients, the resulting granule may also be coated with the enteric material.

### ***Diagnosis of IBS in mammals***

The guidelines for IBS diagnosis promulgated under the Rome criteria are focused upon subjective symptoms, including the continuous or recurrent symptoms of abdominal pain or discomfort that may be relieved with defecation, a change in frequency or

consistency of stools, and that, at least 25 percent of the time, the patient experiences altered stool frequency (greater than 3 bowel movement per day or less than 3 bowel movements per week), altered stool form (hard or loose watery stools or poorly formed stools), passage of mucous stools, or bloating (feeling of abdominal distention). The present invention provides a method of diagnosing irritable bowel syndrome in a patient, based on the patient's malabsorption of carbohydrates. A patient that has one or more IBS symptoms (*e.g.*, a symptom classified under the Rome Criteria) is identified, and a biological sample is obtained from this identified patient. The biological sample can be, *e.g.*, fecal material, urine, blood, serum, plasma, or breath. The amount of a product of a gastrointestinal enzyme in the patient-derived sample is then determined. For example, the hydrogen gas breath test is used to measure hydrogen gas, which is produced as a result of breakdown of unabsorbed carbohydrates in the gastrointestinal tract. The amount of the product in the patient-derived sample is compared with a reference amount of a product of a gastrointestinal enzyme. This reference amount is obtained from a patient or plurality of patients known not to have IBS or other disorders involving carbohydrate malabsorption. An alteration in the test amount relative to the reference amount indicates that the patient has irritable bowel syndrome. For example, an increase in hydrogen gas in a patient-derived sample as compared to a reference sample, measured as described above, indicates poor carbohydrate absorption in the gastrointestinal tract of the patient, leading to the diagnosis of IBS.

## 20 ***Therapeutic Administration***

A therapeutic regimen is carried out by identifying a subject, *e.g.*, a human patient suffering from (or at risk of developing) IBS and providing treatment to the subject. For example, patients characterized as producing less than normal amounts of enzymes that degrade carbohydrates, *e.g.* lactase or fructase, or other digestive enzymes such as amylases, lipases or peptidases, are diagnosed as suffering from or at risk of developing IBS, as now described. A composition including *Bacillus coagulans* bacteria is administered to the patient, such as by oral administration, such that one symptom of IBS is reduced. In embodiments of the invention, the patient is a female, such as a post-menstrual female, since symptoms of IBS are often more prevalent and/or severe in post-menstrual women. The patient may be a post-menopausal woman. The composition including *Bacillus coagulans* bacteria is provided prior to or concomitant with the onset of one or more symptoms of IBS.

Prior to the present invention, dietary management of IBS and other diseases associated with carbohydrate malabsorption has focused on the dietary control of a patient's intake of carbohydrate. The present invention provides a method of dietary management of a subject's carbohydrate absorption, by identifying a patient having a symptom of carbohydrate malabsorption, and providing *Bacillus coagulans* bacteria to said subject, which then colonize said subject's gastrointestinal tract, and cause the subject's carbohydrate absorption to be modulated. Generally, the colonization of *Bacillus coagulans* bacteria in the subject's small and large intestine increases absorption of dietary carbohydrates including fructose and lactose, therefore reducing pain, abdominal bloating, flatus, diarrhea, constipation, and other symptoms of carbohydrate malabsorption.

The present invention also allows the clinician to tailor carbohydrate malabsorption treatment to more effectively manage patient health and wellness. When a patient presents with one or more symptoms of IBS, the physician is able to determine the extent of the patient's carbohydrate malabsorption by measuring the product of a gastrointestinal enzyme such as fructase or lactase in a sample derived from the patient. Measurement of the enzymatic product is performed by methods known in the art and disclosed herein, including the hydrogen and methane breath tests. The physician then provides the compositions of the present invention in amounts appropriate to reduce or eliminate one or more symptoms of IBS.

The present invention further contemplates a therapeutic system for treating, reducing and/or controlling carbohydrate malabsorption. Typically, the system is in the form of a package containing a therapeutic composition of the present invention, or in combination with packaging material. The packaging material includes a label or instructions for use of the components of the package. The instructions indicate the contemplated use of the packaged component as described herein for the methods or compositions of the invention.

By way of example, and not of limitation, a system can comprise one or more unit dosages of a composition according to the present invention. Alternatively, the system can alternately contain bulk quantities of a composition. The label contains instructions for using the therapeutic composition in either unit dose or in bulk forms as appropriate, and may also include information regarding storage of the composition, disease indications, dosages, routes and modes of administration and the like information. Furthermore, depending upon

the particular contemplated use, the system may optionally contain either combined or in separate packages one or more of the following components: bifidogenic oligosaccharides, flavorings, carriers, and the like components. One particularly preferred embodiment comprises unit dose packages of *Bacillus coagulans* bacteria, for use in combination with a conventional liquid product, together with instructions for combining the bacteria with the liquid product, for use in a therapeutic method.

The methods and compositions described herein are useful in the treatment of Celiac disease. Celiac disease is a hereditary disorder that is caused by sensitivity to the gliadin fraction of gluten, a cereal protein found in wheat and rye and less so in barley and oats. The prevalence of celiac disease varies from about 1:300 in southwest Ireland to about 1:5000 in North America. No single genetic marker exists. Celiac disease may be asymptomatic, but most patients have steatorrhea that can range from mild to massive. Symptoms are usually absent until food containing gluten has been eaten. The subject then begins to pass pale, malodorous, bulky stools, and suffers painful abdominal bloating. Thus, a diagnosis is made on the basis of the symptoms and signs, enhanced by laboratory and x-ray studies, and confirmed by biopsy showing a flat mucosa and by subsequent clinical and histologic improvement on a gluten-free diet. Also, the 5-g D-xylose test is usually abnormal, and untreated patients have low C3 and C4, which rise with gluten withdrawal, and normal or increased serum IgA; in 33% to 50%, IgM is reduced.

The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

## EXAMPLES

### EXAMPLE 1. Preparation of *Bacillus coagulans*

#### I. Preparation of Vegetative *Bacillus coagulans*

*Bacillus coagulans* is aerobic and facultative, and is typically cultured at pH 5.7 to 6.8, in a nutrient broth containing up to 2% (by wt) NaCl, although neither NaCl, nor KCl are required for growth. A pH of about 4.0 to about 7.5 is optimum for initiation of sporulation (i.e., the formation of spores). The bacteria are optimally grown at 20°C to 45°C, and the spores can withstand pasteurization. Additionally, the bacteria exhibit facultative and

heterotrophic growth by utilizing a nitrate or sulfate source. The metabolic characteristics of *Bacillus coagulans* are summarized in FIG. 1. However, *Bacillus coagulans* strains and their growth requirements have been described previously (see *e.g.*, Baker, D. *et al*, 1960. *Can. J. Microbiol.* 6: 557-563; Nakamura, H. *et al*, 1988. *Int. J. Syst. Bacteriol.* 38: 63-73. In addition, various strains of *Bacillus coagulans* can also be isolated from natural sources (*e.g.*, heat-treated soil samples) using well-known procedures (see *e.g.*, *Bergey's Manual of Systemic Bacteriology*, Vol. 2, p. 1117, Sneath, P.H.A. *et al.*, eds., Williams & Wilkins, Baltimore, MD, 1986).

*Bacillus coagulans* can be cultured in a variety of media, although it has been demonstrated that certain growth conditions are more efficacious at producing a culture which yields a high level of sporulation. For example, sporulation is demonstrated to be enhanced if the culture medium includes 10 mg/l of MgSO<sub>4</sub> sulfate, yielding a ratio of spores to vegetative cells of approximately 80:20. In addition, certain culture conditions produce a bacterial spore which contains a spectrum of metabolic enzymes particularly suited for the present invention (*i.e.*, production of lactic acid and enzymes for the enhanced probiotic activity and biodegradation). Although the spores produced by these aforementioned culture conditions are preferred, various other compatible culture conditions which produce viable *Bacillus coagulans* spores may be utilized in the practice of the present invention.

Suitable media for the culture of *Bacillus coagulans* include: TSB (Tryptic Soy Broth), GYE (Glucose Yeast Extract Broth), and NB (nutrient broth), which are all well-known within the field and available from a variety of sources. Media supplements which contain enzymatic digests of poultry and/or fish tissue, and containing food yeast are particularly preferred. A preferred supplement produces a media containing at least 60% protein, and about 20% complex carbohydrates and 6% lipids. Media can be obtained from a variety of commercial sources, notably DIFCO (Newark, NJ); BBL (Cockeysville, MD); and Troy Biologicals (Troy, MD).

## II. Preparation of *Bacillus coagulans* Spores

Alternately, a culture of dried *Bacillus coagulans* Hammer bacteria (ATCC No. 31284) spores can be prepared as follows. Approximately  $1 \times 10^7$  spores were inoculated into one liter of culture medium containing: 30g (wt./vol.) Tryptic Soy Broth; 10 g of an

enzymatic-digest of poultry and fish tissue; and 10 g MnSO<sub>4</sub>. The culture was maintained for 72 hours under a high oxygen environment at 37<sup>0</sup>C so as to produce a culture having approximately 6x10<sup>9</sup> cells/gram of culture. The culture was then centrifuged to remove the liquid culture medium and the resulting bacterial paste was re-suspended in 100 ml of sterile water and 20% malto-dextrin and lyophilized. The lyophilized bacteria were ground to a fine powder by use of standard good manufacturing practice (GMP) methodologies.

## **EXAMPLE 2. Therapeutic formulations**

### **Formulation #1**

Bacillus coagulans (800 Million CFU)  
10 Manganese Stereate  
Methyl Cellulose  
Blue Lake Dye

### **Formulation #2**

15 Bacillus coagulans (800 Million CFU)  
Fructose (3,000 units)  
Lactose (3,000 units)  
Lipase (1,500 units)  
Manganese Stereate  
20 Methyl Cellulose

### **Formulation #3**

Bacillus coagulans (800 Million CFU)  
Dibasic Calcium Phosphate  
25 Loperamide HCl (2mg)  
Methyl Cellulose  
Manganese Stereate  
Blue Lake Dye

### **Formulation #4**

30 Bacillus coagulans (800 Million CFU)  
Senosides  
Dibasic Calcium Phosphate  
Loperamide HCl (2mg)  
35 Methyl Cellulose  
Manganese Stereate  
Blue Lake Dye

### **Formulation # 5**

40 Lactase 3,000 FCC units  
Fructose 2,000 FCC units  
Micro-Crystalline Cellulose  
Manganese Stearate

Blue Lake Dye #1  
Raspberry Flavor  
Aspartame

5 Formulation #6

Lactase 3,000 FCC units  
Fructose 2,000 FCC units  
Psyllium Husks  
Micro-Crystalline Cellulose

10 Manganese Stearate  
Blue Lake Dye #1

Formulation #7

15 Loperamide 2mg  
Lactase 3,000 FCC units  
Fructose 2,000 FCC units  
Micro-Crystalline Cellulose  
Manganese Stearate  
20 Blue Lake Dye #1

Formulation #8

Senosides  
Lactase 3,000 FCC units  
25 Fructose 2,000 FCC units  
Micro-Crystalline Cellulose  
Manganese Stearate  
Blue Lake Dye #1

30 Formulation #9

Lysine 125mg  
*Bacillus coagulans* ( $1.5 \times 10^{10}$  CFU/g) 53.3mg  
Nu Tab 358mg  
Mannitol 350mg  
35 Red sugar specks 16mg  
Magnesium stearate 8mg  
Flavor-906.300 (Raspberry) 4mg  
FD&C Blue #1 Lake 0.2mg

40 Formulation #10

*Bacillus coagulans* ( $8.0 \times 10^8$  CFU)  
L-lysine  
cellulose  
magnesium Stearate  
45 hydroxypropylmethylcellulose  
maltodextrin  
flavor



FD&C blue lake dye  
red sucrose specks

### **OTHER EMBODIMENTS**

5           Although particular embodiments have been disclosed herein in detail, this has been  
done by way of example for purposes of illustration only, and is not intended to be limiting  
with respect to the scope of the appended claims, which follow. In particular, it is  
contemplated by the inventors that various substitutions, alterations, and modifications may  
be made to the invention without departing from the spirit and scope of the invention as  
10 defined by the claims. The choice of nucleic acid starting material, clone of interest, or  
library type is believed to be a matter of routine for a person of ordinary skill in the art with  
knowledge of the embodiments described herein. Other aspects, advantages, and  
modifications considered to be within the scope of the following claims.

What is claimed is:

1. A method of reducing a symptom of irritable bowel syndrome, comprising identifying a patient suffering from or at risk of developing irritable bowel syndrome, and administering to said patient a composition comprising *Bacillus coagulans* bacteria.
2. The method of claim 1, wherein said bacteria is *Bacillus coagulans* hammer.
3. The method of claim 2, wherein said bacteria is derived from *Bacillus coagulans* hammer strain Accession No. ATCC 31284.
4. The method of claim 1, wherein said composition further comprises a supplementary enzyme, wherein said enzyme is selected from the group consisting of a lactase, a fructase, a lipase, an amylase and a peptidase.
5. The method of claim 1, wherein said symptom is diarrhea or constipation.
6. The method of claim 1, wherein said symptom is alternating diarrhea and constipation.
7. The method of claim 1, wherein said *Bacillus coagulans* bacteria are provided at a concentration of from about  $1 \times 10^8$  to about  $1 \times 10^{10}$  viable bacteria.
8. The method of claim 1, wherein the *Bacillus coagulans* bacteria are in the form of spores.
9. The method of claim 1, wherein the *Bacillus coagulans* bacteria are in the form of vegetative cells.
10. The method of claim 1, wherein said composition further comprises an anti-diarrheal agent.
11. The method of claim 10, wherein said anti-diarrheal agent is selected from the group consisting of loperamide, attapulgit, Croton Lechleri Extract, and calcium polycarbophil.
12. The method of claim 1, wherein said composition further comprises a laxative agent.
13. The method of claim 12, wherein said laxative agent is selected from the group consisting of a senoside, docusate sodium, magnesium hydroxide, and a dietary fiber.

14. A method of reducing a symptom of irritable bowel syndrome, comprising identifying a patient suffering from or at risk of developing irritable bowel syndrome, and administering to said patient a composition comprising an effective IBS inhibiting amount of *Bacillus coagulans* bacteria prior to or concomitant with the onset of one or more IBS symptoms.

15. The method of claim 14, wherein said bacteria is *Bacillus coagulans* hammer.

16. The method of claim 15, wherein said bacteria is derived from *Bacillus coagulans* hammer strain Accession No. ATCC 31284.

17. The method of claim 14, wherein said composition further comprises a supplementary enzyme, wherein said enzyme is selected from the group consisting of a lactase, a fructase, a lipase, an amylase, and a peptidase.

18. The method of claim 14, wherein said *Bacillus coagulans* bacteria are provided at a concentration of from about  $1 \times 10^8$  to about  $1 \times 10^{10}$  viable bacteria.

15. A method of reducing a symptom of irritable bowel syndrome, comprising identifying a patient suffering from or at risk of developing irritable bowel syndrome, and administering to said patient a composition comprising a fructase and a lactase.

16. The method of claim 15, wherein said fructase is provided at a dose of from about 1000IU to about 12,000IU, and wherein said lactase is provided at a dose of from about 1000IU to about 12,000IU.

17. The method of claim 15, wherein said composition further comprises an anti-diarrheal agent.

18. The method of claim 17, wherein said anti-diarrheal agent is selected from the group consisting of loperamide, attapulgit, Croton Lechleri Extract, and calcium polycarbophil.

19. The method of claim 15, wherein said composition further comprises a laxative agent.

20. The method of claim 19, wherein said laxative agent is selected from the group consisting of a senoside, docusate sodium, magnesium hydroxide, and a dietary fiber.

21. A method of diagnosing irritable bowel syndrome in a patient, comprising the steps of:

identifying a patient having a symptom of irritable bowel syndrome;  
providing a patient-derived biological sample from said identified patient;  
determining an amount of a product of a gastrointestinal enzyme in said patient-derived sample; and

comparing said amount in said patient-derived sample with a reference amount of a product of a gastrointestinal enzyme,

whereby an alteration in the test amount relative to the reference amount indicates that said patient has irritable bowel syndrome.

22. The method of claim 21, wherein said symptom is selected from the group consisting of diarrhea, constipation, or alternating diarrhea and constipation.

23. The method of claim 21, wherein said gastrointestinal enzyme is selected from the group consisting of a lactase, a fructase, a lipase and a peptidase.

24. A method of improving stool consistency in a patient afflicted with non-constipated IBS, comprising administering an effective amount of a *Bacillus coagulans* bacteria, wherein said bacteria are provided at a concentration of from about  $1 \times 10^8$  to about  $1 \times 10^{10}$  viable bacteria, wherein stool consistency is improved following said administration.

25. A method of decreasing urgency in a subject afflicted with IBS, comprising administering an effective amount of a *Bacillus coagulans* bacteria, wherein said bacteria are provided at a concentration of from about  $1 \times 10^8$  to about  $1 \times 10^{10}$  viable bacteria wherein urgency is decreased following said administration.

26. A composition comprising a *Bacillus coagulans* bacteria, a supplementary lactase, and a supplementary fructase.

27. The composition of claim 26, wherein said lactase is a  $\beta$ -galactosidase.

28. The composition of claim 26, wherein said lactase is provided in a concentration from about 1000 IU to about 12,000 IU, and wherein said fructase is provided in a concentration from about 1000 IU to about 12,000 IU.

29. The composition of claim 26, wherein said lactase is provided in a concentration of about 3000 IU, and wherein said fructase is provided in a concentration of about 3000 IU.
30. The composition of claim 26, further comprising one or more components selected from the group consisting of an anti-diarrheal agent, a laxative, a vitamin, a mineral, an isolated amino acid, a source of dietary fiber, and an antibiotic.
31. The composition of claim 26, further comprising an isolated amino acid.
32. The composition of claim 30, wherein said component is manganese stearate.
33. The composition of claim 30, wherein said source of dietary fiber is selected from the group consisting of psyllium husk, soy fiber, citrus fiber, beet fiber, pumpkin seed meal, ground flax, black walnut hull, rice fiber, hydrocollodial polysaccharides, pecan husks, and peanut husks.
34. The composition of claim 26, further comprising a pharmaceutically-acceptable carrier, wherein said carrier comprises silicone.
35. The composition of claim 26, wherein said *Bacillus coagulans* bacteria is provided at a concentration of from about  $1 \times 10^8$  to about  $1 \times 10^{10}$  viable bacteria.
36. The composition of claim 26, wherein said composition is in the form of a capsule, tablet, powder, or liquid.
37. The method of claim 26, wherein said *Bacillus coagulans* bacteria is derived from *Bacillus coagulans* hammer strain Accession No. ATCC 31284.
38. A composition comprising from about  $1 \times 10^8$  to about  $1 \times 10^{10}$  *Bacillus coagulans* bacteria, a supplemental lactase provided in a concentration of about 3000 IU, a supplemental fructase provided in a concentration of about 3000 IU, and manganese stearate.
39. A composition comprising an isolated lactase and an isolated fructase, wherein said isolated lactase is provided in a concentration from about 1000 IU to about 12,000 IU, and wherein said isolated fructase is provided in a concentration from about 1000 IU to about 12,000 IU.

40. The composition of claim 39, further comprising one or more components selected from the group consisting of an anti-diarrheal agent, a laxative, a vitamin, a mineral, an isolated amino acid, a source of dietary fiber, and an antibiotic.
41. The method of claim 40, wherein said anti-diarrheal agent is selected from the group consisting of loperamide, attapulgit, Croton Lechleri Extract, and calcium polycarbophil.
42. The method of claim 40, wherein said laxative agent is selected from the group consisting of a senoside, docusate sodium, magnesium hydroxide, and a dietary fiber.
43. A method for increasing carbohydrate absorption in a mammal, comprising administering to a mammal a composition comprising *Bacillus coagulans* bacteria, a supplementary lactase, and a supplementary fructose, wherein carbohydrate absorption is increased following said administration.
44. The method of claim 43, wherein said mammal is diagnosed as suffering from or being at risk of developing a disorder associated with carbohydrate malabsorption.
45. The method of claim 44, wherein said disorder associated with carbohydrate malabsorption is selected from the group consisting of: lactose intolerance, fructose intolerance, glucose-galactose intolerance, sorbitol intolerance, irritable bowel syndrome, short bowel syndrome, stagnant loop syndrome, celiac disease, chronic malnutrition, chronic persistent diarrhea, immunoproliferative small intestinal disease, intractable diarrhea of infancy, postenteritis syndrome, tropical sprue, Whipple's disease, Wolman disease, Crohn's disease and ulcerative colitis.
46. The method of claim 43, wherein the mammal is human.
47. The method of claim 43, wherein said lactase is provided in a concentration from about 1000 IU to about 12,000 IU, and wherein said fructose is provided in a concentration from about 1000 IU to about 12,000 IU.
48. The method of claim 43, wherein said composition further comprises one or more components selected from the group consisting of an anti-diarrheal agent, a laxative, a vitamin, a mineral, an isolated amino acid, a source of dietary fiber, and an antibiotic.

49. A method for increasing lactose digestion, comprising identifying a patient suffering from or at risk of developing lactose intolerance, and administering to said patient a composition comprising *Bacillus coagulans* bacteria and a supplemental lactase, whereby lactose digestion is increased following said administration.
50. The method of claim 49, wherein the patient is human.
51. The method of claim 49, wherein said lactase is provided in a concentration from about 1000 IU to about 12,000 IU.
52. A composition comprising a *Bacillus coagulans* bacteria and a supplementary fructase.
53. The composition of claim 52, wherein said fructase is provided in a concentration from about 1000 IU to about 12,000 IU.
54. The composition of claim 52, wherein said fructase is provided in a concentration of about 3000 IU.
55. The composition of claim 52, further comprising an isolated amino acid.
56. The composition of claim 52, wherein said composition is in the form of a capsule, tablet, powder, or liquid.
56. The composition of claim 52, wherein said *Bacillus coagulans* bacteria are derived from *Bacillus coagulans* hammer strain Accession No. ATCC 31284.
57. A medical food for the management of irritable bowel syndrome, comprising *Bacillus coagulans* bacteria and L-lysine, wherein said medical food is formulated to provide at least about  $1 \times 10^6$  viable *Bacillus coagulans* bacteria in the gastrointestinal tract of a mammal per day, based on a serving size of about 1 gram to about 2 grams of said medical food taken up to twice a day.
58. The medical food of claim 57, further comprising a supplemental enzyme selected from the group consisting of a lactase, a fructase, a lipase and a peptidase.

59. The medical food of claim 57, further comprising one or more components selected from the group consisting of an anti-diarrheal agent, a laxative, a vitamin, a mineral, an isolated amino acid, a source of dietary fiber, and an antibiotic.

60. A method of dietary management of a subject's carbohydrate absorption, comprising the steps of:

identifying a patient having a symptom of carbohydrate malabsorption; and  
providing a composition comprising *Bacillus coagulans* bacteria to said subject,  
wherein said bacteria colonize said subject's gastrointestinal tract,  
whereby carbohydrate absorption by said subject is modulated, such that the subject's carbohydrate absorption is managed.

61. A method of dietary management of a subject's carbohydrate absorption, comprising the steps of:

identifying a patient having a symptom of carbohydrate malabsorption;  
providing a patient-derived biological sample from said identified patient;  
determining an amount of a product of a gastrointestinal enzyme in said patient-derived sample;  
comparing said amount in said patient-derived sample with a reference amount of a product of a gastrointestinal enzyme; and  
providing a composition comprising *Bacillus coagulans* bacteria,  
whereby the subject's carbohydrate absorption is managed.

62. A method for increasing carbohydrate absorption in a patient diagnosed as suffering from or being at risk of developing celiac disease, comprising administering to said patient a composition comprising *Bacillus coagulans* bacteria, wherein carbohydrate absorption in said patient is increased following said administration.

63. A method of reducing a symptom of irritable bowel syndrome, wherein said symptom comprises alternating diarrhea and constipation, comprising identifying a patient suffering from or at risk of developing irritable bowel syndrome, and administering to said patient a composition comprising *Bacillus coagulans* bacteria.



64. A composition comprising a *Bacillus coagulans* bacterium and a supplementary enzyme provided in a formulation with a food product.

65. The composition of claim 64, wherein said food product is a dairy product.

# **METHODS FOR THE DIETARY MANAGEMENT OF IRRITABLE BOWEL SYNDROME AND CARBOHYDRATE MALABSORPTION**

## **ABSTRACT OF THE INVENTION**

The invention relates generally to digestive disorders, and in particular to methods for treating irritable bowel syndrome by increasing carbohydrate absorption by administering a composition containing a *Bacillus coagulans* bacterium.